

=>


Uploading C:\Program Files\Stnexp\Queries\10698227.str

L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR



G1 C,N

G2 C,O,S,N

Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 15:13:27 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 5696 TO ITERATE

35.1% PROCESSED 2000 ITERATIONS
 INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
 SEARCH TIME: 00.00.01

23 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED ITERATIONS: 109395 TO 118445
 PROJECTED ANSWERS: 825 TO 1795

L2 23 SEA SSS SAM L1

=> s l1 ful

FULL SEARCH INITIATED 15:13:35 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 113496 TO ITERATE

100.0% PROCESSED 113496 ITERATIONS
 SEARCH TIME: 00.00.01

975 ANSWERS

L3 975 SEA SSS FUL L1

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE
 ENTRY

TOTAL
 SESSION

FULL ESTIMATED COST

166.94

167.15

FILE 'CAPLUS' ENTERED AT 15:13:44 ON 16 FEB 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Feb 2006 VOL 144 ISS 8
FILE LAST UPDATED: 15 Feb 2006 (20060215/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s l3

L4 29 L3

=> s l4 and py<2004


23835431 PY<2004

L5 13 L4 AND PY<2004


=> d abs fbib fhitr 1-13

L5 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN

GI



I



II

AB The present invention relates to 3-substituted-2-(aryllalkyl)-1-azabicycloalkanes I [A1 = (CH₂)_n; A2 = (CH₂)_m; A3 = (CH₂)_p; m, n = 1, 2; p = 1 - 4; X = O, NR'; Z = NR', covalent bond, A; A = CR'R'', CR'R''CR'R'', CR':CR', C.tplbond.C (wherein, when Z = bond or A, X = N); Ar = (un)substituted carbocyclic, heterocyclic monocyclic or fused polycyclic aryl; Cy = (un)substituted 5- or 6-membered heteroarom. ring; wavy lines = relative or absolute stereochem. (cis or trans, R or S); R', R'' = H, (un)branched C1-8-alkyl, C3-8-cycloalkyl, heterocyclyl, aryl, aryllalkyl {wherein, substituents = alkyl, alkenyl, heterocyclyl, cycloalkyl, (un)substituted aryl, (un)substituted aryllalkyl, F, Cl, Br, I, OR', NR'R'', CF₃, CN, NO₂, C.tplbond.CR', SR', N₃, C(:O)NR'R'', NR'C(:O)R'', C(:O)R', C(:O)OR', OC(:O)R', O(CR'R'')rC(:O)R', O(CR'R'')rNR''C(:O)R', O(CR'R'')rNR''SO₂R', OC(:O)NR'R'', NR'C(:O)OR'', SO₂R', SO₂NR'R'', NR'SO₂R''}; R'R'' = ring; r = 1 - 6] and II, methods of preparing the compds.

and methods of treatment using the compds. The azabicycloalkanes generally are azabicycloheptanes, azabicyclooctanes, or azabicyclononanes. The aryl group in the arylalkyl moiety is a 5- or 6-membered ring heteroarom., preferably 3-pyridinyl and 5-pyrimidinyl moieties, and the alkyl group is typically a C 1-4 alkyl. The substituent at the 3-position of the 1-azabicycloalkane is a carbonyl group-containing moiety, such as an amide, carbamate, urea, thioamide, thiocarbamate, thiourea or similar functionality. The compds. exhibit activity at nicotinic acetylcholine receptors (nAChRs), particularly the $\alpha 7$ nAChR subtype, and are useful towards modulating neurotransmission and the release of ligands involved in neurotransmission. Methods for preventing or treating conditions and disorders, including central nervous system (CNS) disorders, which are characterized by an alteration in normal neurotransmission, are also disclosed. Also disclosed are methods for treating inflammation, autoimmune disorders, pain and excess neovascularization, such as that associated with tumor growth.

AN 2004:3665 CAPLUS

DN 140:77298

TI Preparation of 3-substituted-2(arylalkyl)-1-azabicycloalkanes and methods of treatment using these compounds

IN Mazurov, Anatoly A.; Klucik, Jozef; Miao, Lan; Seamans, Angela S.; Phillips, Teresa Youngpeter; Schmitt, Jeffrey Daniel; Miller, Craig Harrison

PA Targacept, Inc., USA

SO U.S. Pat. Appl. Publ., 38 pp., Cont.-in-part of U.S. Ser. No. 162,129. CODEN: USXXCO

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2004002513	A1	20040101	US 2003-372642	20030221
	US 6953855	B2	20051011		
				US 1998-210113	A1 19981211
				US 2002-162129	A2 20020604
	US 6432975	B1	20020813	US 1998-210113	19981211 <--
	US 2003045523	A1	20030306	US 2002-162129	20020604 <--
				US 1998-210113	A1 19981211
	CA 2514135	AA	20040910	CA 2004-2514135	20040220
				US 2003-372642	A 20030221
				WO 2004-US5044	W 20040220
	WO 2004076449	A2	20040910	WO 2004-US5044	20040220
	WO 2004076449	A3	20041216		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI				
	RW:				
	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
				US 2003-372642	A 20030221
EP 1594869		A2	20051116	EP 2004-713356	20040220
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
				US 2003-372642	A 20030221
				WO 2004-US5044	W 20040220
US 2005255040		A1	20051117	US 2005-157119	20050620
				US 1998-210113	A1 19981211

			US 2002-162129	B2 20020604
			US 2003-372642	A1 20030221
NO 2005004052	A	20051021	NO 2005-4052	20050831
			US 2003-372642	A 20030221
			WO 2004-US5044	W 20040220

PATENT FAMILY INFORMATION:

FAN 2000:401824

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000034276	A1	20000615	WO 1999-US19906	19990831
	W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW				
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	US 6432975	B1	20020813	US 1998-210113	A 19981211
	CA 2317025	AA	20000615	US 1998-210113	19981211
				CA 1999-2317025	19990831
				US 1998-210113	A 19981211
				WO 1999-US19906	W 19990831
	AU 9956993	A1	20000626	AU 1999-56993	19990831
				US 1998-210113	A 19981211
				WO 1999-US19906	W 19990831
	BR 9907169	A	20001017	BR 1999-7169	19990831
				US 1998-210113	A 19981211
				WO 1999-US19906	W 19990831
	EP 1053236	A1	20001122	EP 1999-944011	19990831
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
				US 1998-210113	A 19981211
				WO 1999-US19906	W 19990831
	JP 2002531564	T2	20020924	JP 2000-586722	19990831
				US 1998-210113	A 19981211
				WO 1999-US19906	W 19990831

OS MARPAT 140:77298

IT 639489-48-8P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of 3-substituted-2(arylalkyl)-1-azabicycloalkanes exhibiting activity at nicotinic acetylcholine receptors)


RN 639489-48-8 CAPLUS

CN 1H-Benzimidazole-5-carboxamide, 1-(1-methylethyl)-N-[(2R,3R)-2-(3-pyridinylmethyl)-1-azabicyclo[2.2.2]oct-3-yl]-2-(trifluoromethyl)- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



L5 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB N-(azabicyclic)arylamides, such as RNR1C(:X)W [R = azabicyclic; R1 = H, alkyl, cycloalkyl, haloalkyl, aryl; W = heteroaryl; X = O, S], were prepared for therapeutic use as nicotinic acetylcholine receptor agonists. These amides are useful for the treatment of central nervous system disorders, such as cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis, attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with

Lewy

Bodies, Huntington's disease, depression, general anxiety disorder, age-related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma, or symptoms associated with pain. Thus, the hydrochloride salt of amide I was prepared via a multistep synthetic sequence which concluded with an amidation reaction of pyrrolo[1,2-c]pyrimidine-3-carboxylic acid hydrochloride with (R)-(+)-3-aminoquinuclidine dihydrochloride using diphenylphosphinic chloride and Et3N in THF. The prepared amides were assayed for human $\alpha 7$ -5HT3 receptor binding activity.

AN 2003:678814 CAPLUS
 DN 139:214613
 TI Preparation of N-(azabicyclyl)arylamides for therapeutic use as nicotinic
 acetylcholine receptor agonists
 IN Rogers, Bruce N.; Piotrowski, David W.; Walker, Daniel P.; Jacobsen, Eric
 Jon; Acker, Brad A.; Wishka, Donn G.; Groppi, Vincent E., Jr.
 PA Pharmacia & Upjohn Company, USA
 SO PCT Int. Appl., 167 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003070732	A1	20030828	WO 2003-US2687	20030214 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
CA 2476681	AA	20030828	CA 2003-2476681	20030214 <--
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
			WO 2003-US2687	W 20030214
AU 2003219690	A1	20030909	AU 2003-219690	20030214 <--
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
			WO 2003-US2687	W 20030214
US 2003236264	A1	20031225	US 2003-366855	20030214 <--
US 6858613	B2	20050222		
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
EP 1476449	A1	20041117	EP 2003-715958	20030214
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
			WO 2003-US2687	W 20030214
BR 2003007735	A	20050125	BR 2003-7735	20030214
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
			WO 2003-US2687	W 20030214
JP 2005523288	T2	20050804	JP 2003-569639	20030214
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
			WO 2003-US2687	W 20030214
US 2005215584	A1	20050929	US 2004-4365	20041203
			US 2002-357917P	P 20020219
			US 2002-423157P	P 20021101
			US 2003-366855	A1 20030214


OS MARPAT 139:214613
 IT 588720-60-9P

RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(preparation of N-(azabicyclyl)arylamides for therapeutic use as nicotinic acetylcholine receptor agonists)

RN 588720-60-9 CAPLUS


CN Pyrrolo[1,2-a]pyrazine-3-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-6-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB N-(azabicyclyl)arylamides, such as RNR1C(:X)W [R = azabicyclyl; R1 = H, alkyl, cycloalkyl, haloalkyl, aryl; W = heteroaryl; X = O, S], were prepared for therapeutic use as nicotinic acetylcholine receptor agonists. These amides are useful for the treatment of central nervous system disorders, such as cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis, attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with

Lewy Bodies, Huntington's disease, depression, general anxiety disorder, age-related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma, or symptoms associated with pain. Thus, the hydrochloride salt of amide I was

prepared via a multistep synthetic sequence which included an amidation reaction of the corresponding (2S,3R)-azabicyclic amine with 5-benzofurancarboxylic acid. The prepared amides were assayed for human $\alpha 7$ -5HT3 receptor binding activity.

AN 2003:678813 CAPLUS
 DN 139:214612
 TI Preparation of N-(azabicyclic)arylamides for therapeutic use as nicotinic acetylcholine receptor agonists
 IN Walker, Daniel P.; Piotrowski, David W.; Jacobsen, Eric Jon; Acker, Brad A.; Groppi, Vincent E., Jr.
 PA Pharmacia & Upjohn Company, USA
 SO PCT Int. Appl., 145 pp.
 CODEN: PIXXD2
 DT Patent
 LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003070731	A2	20030828	WO 2003-US2682	20030213 <--
	WO 2003070731	A3	20040318		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2476624	AA	20030828	US 2002-357926P	P 20020219
				CA 2003-2476624	20030213 <--
				US 2002-357926P	P 20020219
	AU 2003217275	A1	20030909	WO 2003-US2682	W 20030213
				AU 2003-217275	20030213 <--
				US 2002-357926P	P 20020219
	US 2003232853	A1	20031218	WO 2003-US2682	W 20030213
	US 6894042	B2	20050517	US 2003-366431	20030213 <--
				US 2002-357926P	P 20020219
	EP 1476448	A2	20041117	EP 2003-713317	20030213
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
				US 2002-357926P	P 20020219
	BR 2003007782	A	20050104	WO 2003-US2682	W 20030213
				BR 2003-7782	20030213
				US 2002-357926P	P 20020219
				WO 2003-US2682	W 20030213
	JP 2005523287	T2	20050804	JP 2003-569638	20030213
				US 2002-357926P	P 20020219
				WO 2003-US2682	W 20030213
	US 2004224976	A1	20041111	US 2004-868637	20040615
				US 2002-357926P	P 20020219
				US 2003-366431	A3 20030213
OS	MARPAT 139:214612				
IT	588703-34-8P				
	RL:	PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)			

(preparation of N-(azabicyclyl)arylamides for therapeutic use as nicotinic acetylcholine receptor agonists)


RN 588703-34-8 CAPLUS

CN 5-Benzofurancarboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-3-bromo- (9CI)
(CA INDEX NAME)

Absolute stereochemistry.



L5 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



I

AB N-(azabicyclyl)arylamides, such as RNR1C(:X)W [R = azabicyclyl; R1 = H, alkyl, cycloalkyl, haloalkyl, aryl; W = heteroaryl; X = O, S], were prepared for therapeutic use as nicotinic acetylcholine receptor agonists. These amides are useful for the treatment of central nervous system disorders, such as cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis, attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with

Lewy

Bodies, Huntington's disease, depression, general anxiety disorder, age-related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma, or symptoms associated with pain. Thus, the fumarate salt of amide I was prepared via a multistep synthetic sequence which included intramol. cyclization of trans-3-(tert-butoxycarbonylamino)-4-(2-hydroxyethyl)-1-(phenylmethyl)pyrrolidine to form exo-3-(tert-butoxycarbonylamino)-1-azabicyclo[2.2.1]heptane, which contains the target azabicyclic ring, and subsequent amidation of the corresponding azabicyclic amine with 1,3-benzoxazole-5-carboxylic acid. The prepared amides were assayed for

human $\alpha 7$ -5HT3 receptor binding activity.

AN 2003:356448 CAPLUS

DN 138:368781

TI Preparation of N-(azabicyclyl)arylamides for therapeutic use as nicotinic acetylcholine receptor agonists

IN Walker, Daniel P.; Jacobsen, Eric Jon; Piotrowski, David W.; Wishka, Donn G.; Corbett, Jeffrey W.; Groppi, Vincent E., Jr.; Acker, Brad A.; Rauckhorst, Mark R.

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 116 pp.


CODEN: PIXXD2

DT Patent

LA English


FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003037896	A1	20030508	WO 2002-US31579	20021017 <--
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
			US 2001-344436P	P 20011026
			US 2001-342674P	P 20011221
CA 2464194	AA	20030508	CA 2002-2464194	20021017 <--
			US 2001-344436P	P 20011026
			US 2001-342674P	P 20011221
			WO 2002-US31579	W 20021017
EP 1438308	A1	20040721	EP 2002-784010	20021017
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK			
			US 2001-344436P	P 20011026
			US 2001-342674P	P 20011221
			WO 2002-US31579	W 20021017
BR 2002013760	A	20041019	BR 2002-13760	20021017
			US 2001-344436P	P 20011026
			US 2001-342674P	P 20011221
			WO 2002-US31579	W 20021017
JP 2005511574	T2	20050428	JP 2003-540177	20021017
			US 2001-344436P	P 20011026
			US 2001-342674P	P 20011221
			WO 2002-US31579	W 20021017
OS MARPAT 138:368781				
IT 521278-30-8P				
RL:	PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)			
	(preparation of N-(azabicyclyl)arylamides for therapeutic use as nicotinic acetylcholine receptor agonists)			
RN 521278-30-8 CAPLUS				
CN 5-Benzoxazolecarboxamide, N-(6-methyl-1-azabicyclo[2.2.2]oct-3-yl) - (9CI)				
	(CA INDEX NAME)			



RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



II

AB N-(azabicyclyl)arylamides, such as RNR1C(:X)W [R = azabicyclyl; R1 = H, alkyl, cycloalkyl, haloalkyl, aryl; R2 = H, benzyl, alkyl, haloalkyl, cycloalkyl, aryl; W = heteroaryl; X = O, S], were prepared for therapeutic use as nicotinic acetylcholine receptor agonists. These amides are useful for the treatment of central nervous system disorders, such as cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis, attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with Lewy Bodies, Huntington's disease, depression, general anxiety disorder, age-related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma, or symptoms associated with pain. Thus, the fumarate salt of amide II was prepared via a multistep synthetic sequence which included intramol. cyclization of trans-3-(tert-butoxycarbonylamino)-4-(2-hydroxyethyl)-1-phenylmethylpyrrolidine to form exo-3-(tert-butoxycarbonylamino)-1-azabicyclo[2.2.1]heptane, which contains the target azabicyclic ring, and subsequent amidation of the the corresponding azabicyclic amine with furo[2,3-b]pyridine-5-carboxylic acid. The prepared amides were assayed for human $\alpha 7$ -5HT3 receptor binding activity.

AN 2003:282570 CAPLUS

DN 138:304175

TI Preparation of N-(azabicyclyl)arylamides for therapeutic use as nicotinic acetylcholine receptor agonists

IN Walker, Daniel Patrick; Piotrowski, David W.; Jacobsen, Eric Jon; Acker, Brad A.; Wishka, Donn G.; Reitz, Steven Charles; Groppi, Vincent E., Jr.

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 200 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2003029252	A1	20030410	WO 2002-US29827	20021001 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
			US 2001-326565P	P 20011002
			US 2001-326629P	P 20011002
			US 2001-334886P	P 20011115
			US 2001-339633P	P 20011212
CA 2462453	AA	20030410	CA 2002-2462453	20021001 <--
			US 2001-326565P	P 20011002
			US 2001-326629P	P 20011002
			US 2001-334886P	P 20011115
			US 2001-339633P	P 20011212
			WO 2002-US29827	W 20021001
US 2003153595	A1	20030814	US 2002-262257	20021001 <--
US 6911543	B2	20050628		
			US 2001-326565P	P 20011002
			US 2001-326629P	P 20011002
			US 2001-334886P	P 20011115
			US 2001-339633P	P 20011212
EP 1432707	A1	20040630	EP 2002-778286	20021001
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
			US 2001-326565P	P 20011002
			US 2001-326629P	P 20011002
			US 2001-334886P	P 20011115
			US 2001-339633P	P 20011212
			WO 2002-US29827	W 20021001
BR 2002013612	A	20040824	BR 2002-13612	20021001
			US 2001-326565P	P 20011002
			US 2001-326629P	P 20011002
			US 2001-334886P	P 20011115
			US 2001-339633P	P 20011212
			WO 2002-US29827	W 20021001
JP 2005508932	T2	20050407	JP 2003-532500	20021001
			US 2001-326565P	P 20011002
			US 2001-326629P	P 20011002
			US 2001-334886P	P 20011115
			US 2001-339633P	P 20011212
			WO 2002-US29827	W 20021001
US 2003176702	A1	20030918	US 2002-272802	20021017 <--
US 6849620	B2	20050201		
			US 2001-334886P	P 20011115
			US 2001-339633P	P 20011212
BG 108650	A	20050430	BG 2004-108650	20040324

			US 2001-326565P	P	20011002
			US 2001-326629P	P	20011002
			US 2001-334886P	P	20011115
			US 2001-339633P	P	20011212
NO 2004001368	A	20040601	NO 2004-1368		20040401
			US 2001-326565P	P	20011002
			US 2001-326629P	P	20011002
			US 2001-334886P	P	20011115
			US 2001-339633P	P	20011212
US 2005222196	A1	20051006	WO 2002-US29827	W	20021001
			US 2005-137912		20050526
			US 2001-326565P	P	20011002
			US 2001-326629P	P	20011002
			US 2001-334886P	P	20011115
			US 2001-339633P	P	20011212
US 2005234092	A1	20051020	US 2002-262257	A1	20021001
			US 2005-139066		20050526
			US 2001-326565P	P	20011002
			US 2001-326629P	P	20011002
			US 2001-334886P	P	20011115
			US 2001-339633P	P	20011212
			US 2002-262257	A3	20021001

OS MARPAT 138:304175


IT 508208-04-6P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of N-(azabicycyl)arylamides for therapeutic use as nicotinic acetylcholine receptor agonists)



RN 508208-04-6 CAPLUS

CN Furo[2,3-c]pyridine-5-carboxamide, N-(6-methyl-1-azabicyclo[2.2.2]oct-3-yl)- (9CI) (CA INDEX NAME)



RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB 7-Aza[2.2.1]bicycloheptane derivs., such as amides I [R1 = H, alkyl, cycloalkyl, haloalkyl, aryl; R2 = H, benzyl, alkyl, haloalkyl, cycloalkyl, aryl; W = heteroaryl; X = O, S], were prepared for therapeutic use as nicotinic acetylcholine receptor agonists. These amides are useful for the treatment of central nervous system disorders, such as cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis, attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with Lewy Bodies, Huntington's disease, depression, general anxiety disorder, age-related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma, or symptoms associated with pain. Thus, amide dihydrochloride II was prepared via a multistep synthetic sequence which included cycloaddn. of N-tert-butoxycarbonylpyrrole with BrC.tplbond.CCO2Me to form the azabicyclic ring, and subsequent amidation reaction of tert-Bu (1S,2R,4R)-2-amino-7-azabicyclo[2.2.1]heptane-7-carboxylate with 3-methylfuro[2,3-c]pyridine-5-carboxylic acid. The prepared amides were assayed for human α 7-5HT3 receptor binding activity.

AN 2003:221697 CAPLUS

DN 138:238006

TI Preparation of N-[7-aza[2.2.1]bicycloheptanyl]arylamides for therapeutic use as nicotinic acetylcholine receptor agonists

IN Wishka, Donn G.; Walker, Daniel Patrick; Corbett, Jeffrey W.; Reitz, Steven Charles; Rauckhorst, Mark R.; Groppi, Vincent E., Jr.

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 224 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003022856	A1	20030320	WO 2002-US25959	20020904 <--
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
				US 2001-322100P	P 20010912
				US 2001-322333P	P 20010912
				US 2001-322346P	P 20010912
				US 2002-399530P	P 20020730
	CA 2460075	AA	20030320	CA 2002-2460075	20020904 <--
				US 2001-322100P	P 20010912
				US 2001-322333P	P 20010912

			US 2001-322346P	P	20010912
			US 2002-399530P	P	20020730
			WO 2002-US25959	W	20020904
US 2003105089	A1	20030605	US 2002-234575		20020904 <--
			US 2001-322100P	P	20010912
			US 2001-322333P	P	20010912
			US 2001-322346P	P	20010912
			US 2002-399530P	P	20020730
EP 1425286	A1	20040609	EP 2002-757132		20020904
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
			US 2001-322100P	P	20010912
			US 2001-322333P	P	20010912
			US 2001-322346P	P	20010912
			US 2002-399530P	P	20020730
			WO 2002-US25959	W	20020904
BR 2002012477	A	20040824	BR 2002-12477		20020904
			US 2001-322100P	P	20010912
			US 2001-322333P	P	20010912
			US 2001-322346P	P	20010912
			US 2002-399530P	P	20020730
			WO 2002-US25959	W	20020904
JP 2005527472	T2	20050915	JP 2003-526930		20020904
			US 2001-322100P	P	20010912
			US 2001-322333P	P	20010912
			US 2001-322346P	P	20010912
			US 2002-399530P	P	20020730
			WO 2002-US25959	W	20020904

OS MARPAT 138:238006

IT 478170-26-2P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of N-[7-aza[2.2.1]bicycloheptanyl]arylamides for therapeutic use as nicotinic acetylcholine receptor agonists)

RN 478170-26-2 CAPLUS


CN Thiazolo[5,4-c]pyridine-6-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB N-quinuclidinyl-heteroaryls, such as amides I [R¹ = H, alkyl, cycloalkyl, haloalkyl, aryl; R² = H, benzyl, alkyl, haloalkyl, cycloalkyl, aryl; W = aryl, heteroaryl; X = O, S], were prepared for therapeutic use in the treatment of central nervous system disorders, such as cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis, attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with Lewy Bodies, Huntington's disease, depression, general anxiety disorder, age-related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma, or symptoms associated with pain. Thus, the fumarate salt of (3R)-N-quinuclidinyl amide II was prepared via the formation of 6-benzoxazolecarboxylic acid in 89% yield by cyclization of 4-amino-3-hydroxybenzoic acid and (MeO)₃C at 100° for 2 h followed by amide formation of the acid with (R)-(+)-3-aminoquinuclidine dihydrochloride using DIEA in a 5:1 mixture of THF/DMF and subsequent fumarate salt formation. The prepared quinuclidine derivs. were assayed for nicotinic acetylcholinergic receptor binding activity using brain cell membrane prepared from male Sprague-Dawley rats.

AN 2002:964354 CAPLUS

DN 138:24866

TI Preparation and formulation of N-quinuclidinyl-heteroaryls as nicotinic acetylcholinergic receptor modulators for the treatment of a variety of central nervous system disorders

IN Walker, Daniel P.; Wishka, Donn G.; Corbett, Jeffrey W.; Rauckhorst, Mark R.; Piotrowski, David W.; Groppi, Vincent E., Jr.

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 101 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002100858	A2	20021219	WO 2002-US16570	20020606 <--
	WO 2002100858	A3	20030220		
	WO 2002100858	C1	20031224		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,			

	PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW		
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG		
	US 2001-297629P	P	20010612
	US 2001-297630P	P	20010612
	US 2001-297631P	P	20010612
	US 2001-297632P	P	20010612
	US 2001-297633P	P	20010612
	US 2001-328548P	P	20011011
	US 2002-373496P	P	20020418
CA 2445471	AA 20021219	CA 2002-2445471	20020606 <--
	US 2001-297629P	P	20010612
	US 2001-297630P	P	20010612
	US 2001-297631P	P	20010612
	US 2001-297632P	P	20010612
	US 2001-297633P	P	20010612
	US 2001-328548P	P	20011011
	US 2002-373496P	P	20020418
	WO 2002-US16570	W	20020606
US 2003073707	A1 20030417	US 2002-163565	20020606 <--
US 6828330	B2 20041207		
	US 2001-297629P	P	20010612
	US 2001-297630P	P	20010612
	US 2001-297631P	P	20010612
	US 2001-297632P	P	20010612
	US 2001-297633P	P	20010612
	US 2001-328548P	P	20011011
	US 2002-373496P	P	20020418
EP 1404674	A2 20040407	EP 2002-778934	20020606
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR		
	US 2001-297629P	P	20010612
	US 2001-297630P	P	20010612
	US 2001-297631P	P	20010612
	US 2001-297632P	P	20010612
	US 2001-297633P	P	20010612
	US 2001-328548P	P	20011011
	US 2002-373496P	P	20020418
	WO 2002-US16570	W	20020606
JP 2004534065	T2 20041111	JP 2003-503625	20020606
	US 2001-297629P	P	20010612
	US 2001-297630P	P	20010612
	US 2001-297631P	P	20010612
	US 2001-297632P	P	20010612
	US 2001-297633P	P	20010612
	US 2001-328548P	P	20011011
	US 2002-373496P	P	20020418
	WO 2002-US16570	W	20020606
US 2004224977	A1 20041111	US 2004-865149	20040610
	US 2001-297629P	P	20010612
	US 2001-297630P	P	20010612
	US 2001-297631P	P	20010612
	US 2001-297632P	P	20010612
	US 2001-297633P	P	20010612
	US 2001-328548P	P	20011011
	US 2002-373496P	P	20020418

US 2002-163565

A3 20020606

OS MARPAT 138:24866

IT 478169-36-7P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation and formulation of N-quinuclidinyl-heteroaryls as nicotinic acetylcholinergic receptor modulators for treatment of a variety of central nervous system disorders)

RN 478169-36-7 CAPLUS


CN 6-Benzoxazolecarboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L5 ANSWER 8 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN

GI



AB N-quinuclidinyl-heteroaryls, such as amides I [R1 = H, alkyl, cycloalkyl, haloalkyl, aryl; R2 = H, benzyl, alkyl, haloalkyl, cycloalkyl, aryl; W = heteroaryl; X = O, S], were prepared for therapeutic use in the treatment of central nervous system disorders, such as cognitive and attention deficit symptoms of Alzheimer's, neurodegeneration associated with diseases such as Alzheimer's disease, pre-senile dementia (mild cognitive impairment), senile dementia, schizophrenia, psychosis, attention deficit disorder, attention deficit hyperactivity disorder, mood and affective disorders, amyotrophic lateral sclerosis, borderline personality disorder, traumatic brain injury, behavioral and cognitive problems associated with brain tumors, AIDS dementia complex, dementia associated with Down's syndrome, dementia associated with Lewy Bodies, Huntington's disease, depression, general anxiety disorder, age-related macular degeneration, Parkinson's disease, tardive dyskinesia, Pick's disease, post traumatic stress disorder, dysregulation of food intake including bulimia and anorexia nervosa, withdrawal symptoms associated with smoking cessation and dependent drug cessation, Gilles de la Tourette's Syndrome, glaucoma, neurodegeneration associated with glaucoma, or symptoms associated with pain. Thus,

(3R)-N-quinuclidinyl amide II was prepared via a multistep synthetic sequence which started from 2-chloro-3-pyridinol and which included intramol. cyclization of 2-chloro-6-(hydroxymethyl)-4-[(trimethylsilyl)ethynyl]-3-pyridinol to form (7-chlorofuro[2,3-c]pyridin-5-yl)methanol in 27% yield using Et₃N in EtOH, elaboration of the alc. to 2,3-dihydrofuro[2,3-c]pyridine-5-carboxylic acid, and, finally, amidation of the acid with (R)-(+)-3-aminoquinuclidine. The prepared quinuclidine derivs. were assayed for nicotinic acetylcholinergic receptor binding activity using brain cell membrane prepared from male Sprague-Dawley rats.

AN 2002:964353 CAPLUS

DN 138:24865

TI Preparation and formulation of N-quinuclidinyl-heteroaryls as nicotinic acetylcholinergic receptor modulators for the treatment of a variety of central nervous system disorders

IN Wishka, Donn G.; Reitz, Steven C.; Piotrowski, David W.; Groppi, Vincent E., Jr.

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 262 pp.

CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002100857	A1	20021219	WO 2002-US16568	20020606 <--
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
				US 2001-297708P	P 20010612
				US 2001-297709P	P 20010612
				US 2001-297710P	P 20010612
				US 2001-297711P	P 20010612
				US 2001-297712P	P 20010612
				US 2001-328596P	P 20011011
				US 2002-373495P	P 20020418
CA 2445467	AA	20021219	CA 2002-2445467	20020606 <--	
			US 2001-297708P	P 20010612	
			US 2001-297709P	P 20010612	
			US 2001-297710P	P 20010612	
			US 2001-297711P	P 20010612	
			US 2001-297712P	P 20010612	
			US 2001-328596P	P 20011011	
			US 2002-373495P	P 20020418	
			WO 2002-US16568	W 20020606	
US 2003045540	A1	20030306	US 2002-163564	20020606 <--	
			US 2001-297708P	P 20010612	
			US 2001-297709P	P 20010612	
			US 2001-297710P	P 20010612	
			US 2001-297711P	P 20010612	
			US 2001-297712P	P 20010612	
			US 2001-328596P	P 20011011	
			US 2002-373495P	P 20020418	

EP 1406901	A1	20040414	EP 2002-778932	20020606
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
			US 2001-297708P	P 20010612
			US 2001-297709P	P 20010612
			US 2001-297710P	P 20010612
			US 2001-297711P	P 20010612
			US 2001-297712P	P 20010612
			US 2001-328596P	P 20011011
			US 2002-373495P	P 20020418
			WO 2002-US16568	W 20020606
BR 2002010384	A	20040629	BR 2002-10384	20020606
			US 2001-297708P	P 20010612
			US 2001-297709P	P 20010612
			US 2001-297710P	P 20010612
			US 2001-297711P	P 20010612
			US 2001-297712P	P 20010612
			US 2001-328596P	P 20011011
			US 2002-373495P	P 20020418
			WO 2002-US16568	W 20020606
CN 1511154	A	20040707	CN 2002-809814	20020606
			US 2001-297708P	P 20010612
			US 2001-297709P	P 20010612
			US 2001-297710P	P 20010612
			US 2001-297711P	P 20010612
			US 2001-297712P	P 20010612
			US 2001-328596P	P 20011011
			US 2002-373495P	P 20020418
JP 2004537532	T2	20041216	JP 2003-503624	20020606
			US 2001-297708P	P 20010612
			US 2001-297709P	P 20010612
			US 2001-297710P	P 20010612
			US 2001-297711P	P 20010612
			US 2001-297712P	P 20010612
			US 2001-328596P	P 20011011
			US 2002-373495P	P 20020418
			WO 2002-US16568	W 20020606
ZA 2003008844	A	20040628	ZA 2003-8844	20031113
			US 2001-297710P	P 20010612
OS	MARPAT 138:24865			
IT	478148-57-1P, N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-7-chlorofuro[2,3-c]pyridine-5-carboxamide hydrochloride			
	RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)			
	(preparation and formulation of N-quinuclidinyl-heteroaryl amides as nicotinic acetylcholinergic receptor modulators for treatment of a variety of central nervous system disorders)			
RN	478148-57-1 CAPLUS			
CN	Furo[2,3-c]pyridine-5-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-7-chloro-, monohydrochloride (9CI) (CA INDEX NAME)			


Absolute stereochemistry.




● HCl

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



I



II

AB The title compds. [I; R = H, OH, amino, halo, CF₃, alkylsulfonyl, sulfamyl; R₁-R₆ = H, alkyl; vicinal pairs of R₁-R₆ - atoms to form 5-7 membered rings, double bonds; R₇ = aminoalkyl, 3- or 4-quinuclidinyl, 4-(1-azabicyclo[3.3.1]nonyl), 4-[3-methoxy-1-(3-[4-fluorophenoxy]propyl)piperidinyl], etc.; R₈ = alkyl, alkylcarbonylalkyl, etc.; n = 1-3], were prepared as CNS gastric prokinetic, and antiemetic agents (no data). Thus, 5-methoxytetralone was refluxed with SnCl₂ in EtOH/conc HCl for 16 h to give 5-methoxytetralin. The latter in DMF was stirred with N-chlorosuccinimide at 0° for 4 h to give 8-chloro-5-methoxytetralin; this was brominated similarly with N-bromosuccinimide to give 6-bromo-8-chloro-5-methoxytetralin. This in THF at -78° was treated with BuLi and then CO₂ to give 8-chloro-5-methoxytetralin-6-carboxylic acid. The acid in CHCl₃/Et₃N at -23° was treated with EtO₂CCl and then aminoquinuclidine dihydrochloride/aqueous K₂CO₃ to give quinuclidinylcarbamoyltetralin II. I are said to lack dopamine D₂ activity.

AN 1990:591177 CAPLUS

DN 113:191177

TI Azabicyclic carbamoylarenes as 5-HT₃ antagonists useful as antiemetics

IN Pelletier, Jeffrey C.; Youssefyeh, Raymond D.; Campbell, Henry F.

PA Rorer Pharmaceutical Corp., USA


SO U.S., 9 pp.

CODEN: USXXAM


DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4920227	A	19900424	US 1988-277611 US 1988-277611	19881129 <-- 19881129


OS MARPAT 113:191177
 IT 129764-46-1P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, as serotonin 5-HT₃ antagonist)
 RN 129764-46-1 CAPLUS
 CN 1H-Indene-5-carboxamide, N-1-azabicyclo[2.2.2]oct-3-yl-7-chloro-2,3-dihydro-4-methoxy- (9CI) (CA INDEX NAME)



L5 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
 GI



I




II

AB The title compds. [I; X = H, OH, amino, alkylamino, halo, CF₃, alkylsulfamyl, alkylsulfonyl, etc.; R = H, alkyl; R₁, R₂ = H, alkyl, vicinal R₂ groups together may be (CH₂)_a; a = 1-4; n = 2-4 V = alkyl, (CR₁R₂)b-SO-R₃, (CR₁R₂)bCOR₃; R₃ = alkyl; Z = (CR₁R₂)d-NR₁R₂, 3- or 4-quinuclidinyl, etc.; b, d = 1-3] and their pharmaceutically acceptable salts, which are 5-HT₃ antagonists and have gastric prokinetic and antiemetic activities and lack D₂ receptor binding activity, were prepared 3-Aminoquinuclidine and a K₂CO₃ solution were added to a mixture of ClCO₂Et and N-acetyl-2,3-dihydroindole II (R₄ = NH₂) in CHCl₃-Et₃N at -20° and the resulting mixture was stirred for 2 h to give II (R₄ = 3-quinuclidinylamino). At 2.0 mg/kg i.v. I showed antiemetic activity in rats treated with cisplatin.



AN 1990:591155 CAPLUS
 DN 113:191155
 TI Preparation of indole, quinoline, and benzazepine analogs as 5-HT₃ antagonists
 IN Pelletier, Jeffrey C.; Youssefyeh, Raymond D.; Campbell, Henry F.
 PA Rorer Pharmaceutical Corp., USA

SO U.S., 15 pp.
CODEN: USXXAM
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4920219	A	19900424	US 1988-277582	19881129 <--
	WO 9006113	A1	19900614	WO 1989-US5422	19891129 <--
	W: AU, JP, US				
	RW: AT, BE, CH, DE, ES, FR, GB, IT, LU, NL, SE				
				US 1988-277582	A1 19881129
	AU 9047439	A1	19900626	AU 1990-47439	19891129 <--
				US 1988-277582	A 19881129
				WO 1989-US5422	A 19891129
	US 5063230	A	19911105	US 1990-489646	19900406 <--
				US 1988-277582	A3 19881129
OS	CASREACT 113:191155; MARPAT 113:191155				
IT	129511-02-0P				
	RL: SPN (Synthetic preparation); PREP (Preparation)				
	(preparation of, as 5-HT3 antagonist)				
RN	129511-02-0 CAPLUS				
CN	1H-Indole-5-carboxamide, N-1-azabicyclo[2.2.2]oct-3-yl-7-chloro-2,3-dihydro-4-methoxy- (9CI) (CA INDEX NAME)				



L5 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB The title compds. [I; 1, m, n, p = 0-3; R1-R4 = H, lower alkyl, Ph; R5 = H, lower alkyl; X = O, S; Ar = (un)substituted Ph, pyridyl, furyl, thienyl, 6-methoxy-1H-benzotriazol-5-yl, 6-methoxy-indol-5-yl, 2-(un)substituted amino-4-methoxypyrimidin-5-yl, 3,4-methylenedioxyphenyl, (un)substituted naphthalenyl, indolyl] useful for the enhancement of memory or the correction of memory deficiency (no data), are prepared. Thus, THF was added to a mixture of 4-amino-5-chloro-2-methoxybenzoic acid and 1,1'-carbonyldimidazole with stirring. When evolution of CO₂ ceased, N was bubbled 1 h through the reaction mixture. A solution of 3-aminoquinuclidine in THF was added to the stirred mixture and stirring at room temperature continued 3 h to give 67% 3-benzamidoquinuclidine derivative (II). A total 46 3-[(hetero)arylamido]quinuclidine including their salts were prepared.

AN 1990:55619 CAPLUS

DN 112:55619

TI Arylamido- and arylthioamidoazabicycloalkanes for enhancing memory or correcting memory deficiency

IN Smith, William Levi

PA A. H. Robins Co., Inc., USA

SO Eur. Pat. Appl., 27 pp.

CODEN: EPXXDW

DT Patent


LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	----	-----	-----	-----
PI	EP 327335	A1	19890809	EP 1989-300961	19890201 <--
	EP 327335	B1	19921014		
	R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	US 4863919	A	19890905	US 1988-150981	A 19880201
	IL 88434	A1	19920906	US 1988-150981	19880201 <--
				IL 1988-88434	19881121 <--
				US 1988-150981	A 19880201
	ZA 8809109	A	19890830	ZA 1988-9109	19881205 <--
				US 1988-150981	A 19880201
	JP 01226818	A2	19890911	JP 1989-13894	19890123 <--
				US 1988-150981	A 19880201
	CA 1320448	A1	19930720	CA 1989-589194	19890126 <--

DK 8900425	A	19890802	US 1988-150981	A	19880201
			DK 1989-425		19890131 <--
AU 8929506	A1	19890803	US 1988-150981	A	19880201
AU 629197	B2	19921001	AU 1989-29506		19890201 <--
AT 81457	E	19921015	US 1988-150981	A	19880201
			AT 1989-300961		19890201 <--
			US 1988-150981	A	19880201
			EP 1989-300961	A	19890201
ES 2045402	T3	19940116	ES 1989-300961		19890201 <--
			US 1988-150981	A	19880201


OS CASREACT 112:55619; MARPAT 112:55619
 IT 106517-99-1P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, for enhancement of memory)
 RN 106517-99-1 CAPLUS
 CN 1H-Indole-5-carboxamide, N-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)



L5 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN
 GI



I



II

AB Carboxylate and sulfonate esters, carboxamides, and sulfonamides of a variety of N-containing heterocyclic alcs. and amines with a variety of mono- and bicyclic carbocyclic and heterocyclic acids and imidazolylmethyltetrahydrocarbazolones I (R1 = H, C1-10 alkyl, C3-9 cycloalkyl, C3-6 alkenyl, Ph, phenylalkyl; R2-R4 = H, C1-6 alkyl, C3-7 cycloalkyl, C2-4 alkenyl, phenylalkyl) were prepared (.apprx.80 compds.) for treatment of psychotic disorders, rhinitis, and pulmonary embolism and to

improve the nasal resorption of other drugs such as peptides.
 endo-8-Methyl-8-azabicyclo[3.2.1]oct-3-yl indole-3-carboxylate (II) at 0.01-100 µg/kg i.p. reversed the stress-induced inhibition of social behavior in mice, and at 1-10 mg/kg orally inhibited the stress-induced elevation of plasma corticosterone in mice in a manner similar to diazepam. II reached a level of 200 ng/mL in the plasma 5-10 mins. after nasal administration, compared to 30-40 mins. after oral administration of the same dose. A nasal spray for treatment of rhinitis or pulmonary embolism contained II-HCl 100 mg, benzalkonium chloride 0.1 mg, 0.9% aqueous NaCl 0.6 mL, and distilled water 0.4 mL. Pseudotropine was chlorinated to 3-chloro-8-methyl-8-azabicyclo[3.2.1]octane, which was converted successively to the 3-cyano, 3-methoxycarbonyl, 3-carboxy, and 3-chlorocarbonyl derivs. followed by reaction with MeMgI and indole to produce 3β-(indole-3-carbonyl)-8-methyl-8-azabicyclo[3.2.1]octane.

AN 1989:8041 CAPLUS

DN 110:8041

TI Preparation and use of carbocyclic and heterocyclic esters and amides and imidazolylcarbazoles for treatment of psychosis, rhinitis, and pulmonary embolism and for facilitation of the nasal resorption of drugs

IN Azria, Moise; Buchheit, Karl Heinz; Dixon, Keith Arnold; Engel, Guenther; Giger, Rudolf Karl Andreas

PA Sandoz-Patent-G.m.b.H., Fed. Rep. Ger.

SO Ger. Offen., 27 pp.

CODEN: GWXXBX

DT Patent

LA German

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 3724059	A1	19880218	DE 1987-3724059	19870721 <--
				GB 1986-18614	A 19860730
				DE 1986-3626703	A1 19860807
	NL 8701682	A	19880216	NL 1987-1682	19870716 <--
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807
	HU 45895	A2	19880928	HU 1987-3252	19870716 <--
	HU 202108	B	19910228		
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807
	FR 2602142	A1	19880205	FR 1987-10519	19870722 <--
	FR 2602142	B1	19960705		
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807
	CH 675072	A	19900831	CH 1987-2831	19870723 <--
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807
	BE 1005921	A4	19940315	BE 1987-818	19870723 <--
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807
	NO 8703133	A	19880201	NO 1987-3133	19870727 <--
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807
	GB 2193633	A1	19880217	GB 1987-17768	19870727 <--
	GB 2193633	B2	19910417		
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807
	DK 8703924	A	19880131	DK 1987-3924	19870728 <--
				GB 1986-18614	A 19860730
				DE 1986-3626703	A 19860807

FI 8703280	A	19880131	FI 1987-3280	19870728 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
AU 8776190	A1	19880204	AU 1987-76190	19870728 <--
AU 610074	B2	19910516		
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
SE 8702980	A	19880428	SE 1987-2980	19870728 <--
SE 504184	C2	19961202		
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
ES 2010227	A6	19891101	ES 1987-2207	19870728 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
IL 83363	A1	19930708	IL 1987-83363	19870728 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
IL 96796	A1	19940731	IL 1987-96796	19870728 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
			IL 1987-83363	A3 19870728
IL 96797	A1	19941229	IL 1987-96797	19870728 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
			IL 1987-83363	A3 19871231
JP 63041429	A2	19880222	JP 1987-193629	19870729 <--
JP 2632858	B2	19970723		
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
AT 8701912	A	19930515	AT 1987-1912	19870729 <--
AT 396870	B	19931227		
			DE 1986-3626703	A 19860807
CA 1327750	A1	19940315	CA 1987-543271	19870729 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
ZA 8705652	A	19890628	ZA 1987-5652	19870730 <--
			GB 1986-18614	A 19860730
ES 2016440	A6	19901101	ES 1989-1137	19890331 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
ZA 8903145	A	19890628	ZA 1989-3145	19890427 <--
			GB 1986-18614	A 19860730
ZA 8903146	A	19890628	ZA 1989-3146	19890730 <--
			GB 1986-18614	A 19860730
GB 2231264	A1	19901114	GB 1990-8068	19900410 <--
GB 2231264	B2	19910424		
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
			GB 1987-17768	A3 19870727
GB 2231265	A1	19901114	GB 1990-8069	19900410 <--
GB 2231265	B2	19910424		
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
			GB 1987-17768	A3 19870727
AU 9171946	A1	19910509	AU 1991-71946	19910227 <--
AU 642210	B2	19931014		
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
AU 9172910	A1	19910516	AU 1991-72910	19910314 <--

AU 637878	B2	19930610	GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
CA 1334075	A1	19950124	CA 1992-616654	19920909 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
			CA 1987-543271	A3 19870729
US 5561149	A	19961001	US 1995-403620	19950314 <--
			GB 1986-18614	A 19860730
			DE 1986-3626703	A 19860807
			US 1987-78336	B1 19870727
			US 1989-423916	B1 19891019
			US 1991-701934	B1 19910517
			US 1992-890493	B1 19920528
			US 1993-3926	B1 19930113
			US 1993-111805	B1 19930825


OS MARPAT 110:8041

IT 117843-77-3P

RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, for lung embolism and mental disorder and rhinitis treatment)


RN 117843-77-3 CAPLUS

CN 1H-Indole-5-carboxamide, N-1-azabicyclo[2.2.2]oct-3-yl-2,3-dihydro- (9CI)
 (CA INDEX NAME)



L5 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2006 ACS on STN

GI



I

AB Title compds. I [R1-R4 = H, alkyl, Ph; R5 = H, alkyl; X = O, S; R6 = (un)substituted Ph, naphthyl; m, o, p = 0-3] and their salts, useful for memory enhancement, were prepared. Thus, 4-amino-5-chloro-2-methoxybenzoic acid in THF was reacted with 1,1'-carbonyldiimidazole and then with 3-aminoquinuclidine to give the free base, which was converted to 4-amino-N-(1-azabicyclo[2.2.2]oct-3-yl)-5-chloro-2-methoxybenzamide-HCl (II). In tests in mice II significantly increased the time required to complete a drinking task, indicating a measure of improved memory.

AN 1987:67141 CAPLUS

DN 106:67141

TI Enhancing memory or correcting memory deficiency with arylamido- and
arylthioamidoazabicycloalkanes

IN Welstead, William J., Jr.

PA A. H. Robins Co., Inc., USA

SO U.S., 12 pp.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4605652	A	19860812	US 1985-697944	19850204 <--
	AU 8652672	A1	19860807	AU 1986-52672	19860123 <--
	AU 589155	B2	19891005		
				US 1985-697944	A 19850204
	CA 1273297	A1	19900828	CA 1986-500990	19860203 <--
				US 1985-697944	A 19850204
	EP 190920	A2	19860813	EP 1986-300747	19860204 <--
	EP 190920	A3	19900418		
	EP 190920	B1	19930505		
	R: AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE				
				US 1985-697944	A 19850204
	JP 61183223	A2	19860815	JP 1986-22819	19860204 <--
	JP 06062414	B4	19940817		
				US 1985-697944	A 19850204
	AT 88891	E	19930515	AT 1986-300747	19860204 <--
				US 1985-697944	A 19850204
				EP 1986-300747	A 19860204


OS MARPAT 106:67141

IT 106517-99-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, for memory improvement)


RN 106517-99-1 CAPLUS

CN 1H-Indole-5-carboxamide, N-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX
NAME)



=> d his

L6 ANSWER 12 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



II

AB The invention discloses compds. that are selective $\alpha 7$ nAChR agonists and 5-HT₃ antagonists (no data), specifically, compds. Azb-NH-CO-W0 [I; Azb = selected azabicycloalkyl, notably 1-azabicyclo[2.2.2]oct-3-yl, 1-azabicyclo[2.2.1]hept-3-yl, exo-7-azabicyclo[2.2.1]hept-2-yl, and 1-azabicyclo[3.2.1]oct-3-yl, all with optional alkyl or substituted alkyl substituents or N-protecting groups; W0 = selected 6/5-bicyclic heteroaryls, notably thieno[3,2-c]pyridine, 1-benzofuran, pyrrolo[2,3-c]pyridine, furo[2,3-c]pyridine, thieno[2,3-c]pyridine, and pyrrolo[1,2-a]pyrazine]. Compds. I are useful for treating many CNS diseases, including schizophrenia, psychosis, Alzheimer's and other neurodegenerative diseases, emesis, migraine, anxiety, and substance dependence withdrawal. Approx. 20 synthetic examples, some with data for the products, are given, as well as prepsns. of various azabicycloalkylamine and heterobicycloarom. carboxylic acid precursors. For instance, thieno[3,2-c]pyridine-6-carboxylic acid was prepared from glyoxylic acid and 2,3-thiophenedicarboxaldehyde in approx. 6 steps. Amidation of this acid with (R)-3-aminoquinuclidine using bis(2-oxo-3-oxazolidinyl)phosphinic chloride and TEA gave invention compound II, isolated as the di-HCl salt.

AN 2004:390256 CAPLUS

DN 140:406824

TI Compounds having both $\alpha 7$ nicotinic agonist activity and 5-HT₃ antagonist activity, for treatment of CNS diseases, and their preparation, pharmaceutical compositions, and use

IN Wong, Erik Ho Fong; Cortes-Burgos, Luz Amparo; Rogers, Bruce Nelsen; Piotrowski, David Walter; Walker, Daniel Patrick; Jacobsen, Eric Jon; Wishka, Donn Gregory; Acker, Brad Alan

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 77 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004039815	A2	20040513	WO 2003-IB4681	20031020
	WO 2004039815	A3	20040722		
	WO 2004039815	C1	20040923		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,				

TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,
 FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

CA 2503786	AA	20040513	CA 2003-2503786	20031020
BR 2003015056	A	20050816	BR 2003-15056	20031020
EP 1562959	A2	20050817	EP 2003-751183	20031020

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK

US 2004147522	A1	20040729	US 2003-698227	20031031
---------------	----	----------	----------------	----------


PRAI US 2002-423155P P 20021101
 WO 2003-IB4681 W 20031020

OS MARPAT 140:406824
 IT 588720-60-9P

RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (drug candidate; preparation of azabicycloalkyl heterobicycloarenecarboxamides as $\alpha 7$ nicotinic agonists and 5-HT3 antagonists)


RN 588720-60-9 CAPLUS
 CN Pyrrolo[1,2-a]pyrazine-3-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-6-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



after priority
date

L6 ANSWER 10 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB Title N-(1-azabicyclo[2.2.2]octyl)heteroarylamides I and analogs [wherein X = o, S; R¹ = H, (halo)alkyl, cycloalkyl, substituted Ph, naphthyl; R² = independently halo, cycloalkyl, aryl, (un)substituted alkyl; m = 0-1; n = 0-1; with the proviso that m + n = 1; W = (un)substituted Ph, heterocyclyl, heteroaryl; or pharmaceutically acceptable salts, racemic mixts., or pure enantiomers thereof] were prepared as $\alpha 7$ nicotinic acetylcholine receptor (nAChR) full agonists (no data). For example, reaction of phosgene with 4-bromopyrazole in EtOAc, followed by coupling with (+)-3-aminoquinuclidine•2HCl provided II•HCl (25%). The invention provides for compns. of I with psychostimulants and/or monoamine reuptake inhibitors for the treatment of attention deficit hyperactivity disorder (ADHD).

AN 2004:513575 CAPLUS

DN 141:71755

TI Preparation of N-(quinuclidinyl)heteroarylamides as nicotinic acetylcholine receptor agonists for use in combination therapy for the treatment of ADHD

IN Groppi, Vincent Edward, Jr.; Jacobsen, Eric Jon; Myers, Jason Kenneth; Piotrowski, David Walter; Rogers, Bruce Nelsen; Walker, Daniel Patrick; Wishka, Donn Gregory

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 141 pp.

CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004052461	A1	20040624	WO 2003-IB5542	20031128
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2509142	AA	20040624	CA 2003-2509142	20031128
	EP 1572300	A1	20050914	EP 2003-775637	20031128
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
	BR 2003017229	A	20051101	BR 2003-17229	20031128

US 2005107425 A1 20050519 US 2004-963922 20041012
 NO 2005003185 A 20050817 NO 2005-3185 20050629
 PRAI US 2002-432586P P 20021211
 WO 2003-IB5542 W 20031128
 US 2003-731402 B1 20031209
 OS MARPAT 141:71755
 IT 478148-80-0P, N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-3-methylfuro[2,3-c]pyridine-5-carboxamide
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (nAChR agonist; preparation of N-(quinuclidinyl)heteroarylamides as nAChR agonists for use in combination therapy for treatment of ADHD)
 RN 478148-80-0 CAPLUS
 CN Furo[2,3-c]pyridine-5-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-3-methyl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 11 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to azabicycloalkane derivs. of formula azabicyclo-N(R1)-C(:X)-W [wherein: R1 is H, (cyclo)alkyl, or haloalkyl, etc.; X is O or S; W is a substituted benzene], useful as $\alpha 7$ nAChR agonists. Pharmacokinetics of the prepared compds. were evaluated (no biol. data). Blood-brain barrier penetration was investigated (no biol. data). For instance, chiral azabicycloheptane derivative I was prepared via addition

of Me

3-bromopropargylate to N-Boc-pyrrole, reduction of the obtained azabicyclo[2.2.1]heptadiene II, hydrolysis of the obtained azabicycloheptane derivative III (R2 = OMe), reaction of the carboxylic acid III (R2 = OH) with diphenylphosphoryl azide and benzyl alc., resolution of the obtained exo-derivative IV, and hydrogenation.

AN 2004:513522 CAPLUS
 DN 141:71300

TI A preparation of azabicycloalkane derivatives, useful as $\alpha 7$ nicotinic acetylcholine receptor ($\alpha 7$ nAChR) agonists

IN Corbett, Jeffrey Wayne; Groppi, Vincent Edward, Jr.

PA Upjohn Company, USA

SO PCT Int. Appl., 151 pp.


CODEN: PIXXD2

DT Patent


LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004052348	A2	20040624	WO 2003-IB5525	20031128
	WO 2004052348	A3	20041021		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2508004	AA	20040624	CA 2003-2508004	20031128
	EP 1572205	A2	20050914	EP 2003-772599	20031128
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
	BR 2003017110	A	20051025	BR 2003-17110	20031128
	US 2005245504	A1	20051103	US 2003-731565	20031209
PRAI	US 2002-432527P	P	20021211		
	WO 2003-IB5525	W	20031128		
OS	MARPAT 141:71300				
IT	478148-80-0P				
	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(preparation of azabicycloalkane derivs. useful as $\alpha 7$ nAChR agonists)				
RN	478148-80-0 CAPLUS				
CN	Furo[2,3-c]pyridine-5-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-3-methyl- (9CI) (CA INDEX NAME)				

Absolute stereochemistry.



L6 ANSWER 12 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB The invention discloses compds. that are selective $\alpha 7$ nAChR agonists and 5-HT₃ antagonists (no data), specifically, compds. Azb-NH-CO-W0 [I; Azb = selected azabicycloalkyl, notably 1-azabicyclo[2.2.2]oct-3-yl, 1-azabicyclo[2.2.1]hept-3-yl, exo-7-azabicyclo[2.2.1]hept-2-yl, and 1-azabicyclo[3.2.1]oct-3-yl, all with optional alkyl or substituted alkyl substituents or N-protecting groups; W0 = selected 6/5-bicyclic heteroaryls, notably thieno[3,2-c]pyridine, 1-benzofuran, pyrrolo[2,3-c]pyridine, furo[2,3-c]pyridine, thieno[2,3-c]pyridine, and pyrrolo[1,2-a]pyrazine]. Compds. I are useful for treating many CNS diseases, including schizophrenia, psychosis, Alzheimer's and other neurodegenerative diseases, emesis, migraine, anxiety, and substance dependence withdrawal. Approx. 20 synthetic examples, some with data for the products, are given, as well as prepsns. of various azabicycloalkylamine and heterobicycloarom. carboxylic acid precursors. For instance, thieno[3,2-c]pyridine-6-carboxylic acid was prepared from glyoxylic acid and 2,3-thiophenedicarboxaldehyde in approx. 6 steps. Amidation of this acid with (R)-3-aminoquinuclidine using bis(2-oxo-3-oxazolidinyl)phosphinic chloride and TEA gave invention compound II, isolated as the di-HCl salt.

AN 2004:390256 CAPLUS

DN 140:406824

TI Compounds having both $\alpha 7$ nicotinic agonist activity and 5-HT₃ antagonist activity, for treatment of CNS diseases, and their preparation, pharmaceutical compositions, and use

IN Wong, Erik Ho Fong; Cortes-Burgos, Luz Amparo; Rogers, Bruce Nelsen; Piotrowski, David Walter; Walker, Daniel Patrick; Jacobsen, Eric Jon; Wishka, Donn Gregory; Acker, Brad Alan

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 77 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004039815	A2	20040513	WO 2003-IB4681	20031020
	WO 2004039815	A3	20040722		
	WO 2004039815	C1	20040923		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2503786	AA	20040513	CA 2003-2503786	20031020
	BR 2003015056	A	20050816	BR 2003-15056	20031020
	EP 1562959	A2	20050817	EP 2003-751183	20031020
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	US 2004147522	A1	20040729	US 2003-698227	20031031
PRAI	US 2002-423155P	P	20021101		
	WO 2003-IB4681	W	20031020		
OS	MARPAT 140:406824				
IT	588720-60-9P				


application

RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (drug candidate; preparation of azabicycloalkyl heterobicycloarenecarboxamides as $\alpha 7$ nicotinic agonists and 5-HT3 antagonists)


RN 588720-60-9 CAPLUS

CN Pyrrolo[1,2-a]pyrazine-3-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-6-bromo- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 13 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 GI



AB Quinuclidine derivs., such as $RNHC(:X)W$, $RC(:X)NHW$, $RNHCH_2W$ and RCH_2NHW [R = quinuclidinyl; W = indazolyl, benzothiazolyl, benzoisothiazolyl; X = O, S], were prepared for therapeutic use as nicotinic acetylcholine receptor $\alpha 7$ ($\alpha 7$ nAChR) ligands for the treatment of psychotic or neurodegenerative diseases and disorders involving dysfunction of the cholinergic system. These quinuclidines are claimed for use in the treatment of dementia or memory impairment due to mild cognitive impairment due to Alzheimer's disease, schizophrenia, Parkinson's disease, Huntington's disease, Pick's disease, Creutzfeldt-Jakob disease, depression, aging, head trauma, stroke, CNS hypoxia, cerebral senility, or multiinfarct dementia. These quinuclidines are also claimed for use in the treatment of intoxication, damage associated with strokes, ischemia and glutamate-induced excitotoxicity, smoking cessation or nicotine addiction, pain, jet lag, obesity, diabetes, mild cognitive impairment (MCI), vascular dementia (VaD), age-associated cognitive decline (AACD), amnesia associated with open-heart-surgery, cardiac arrest, general anesthesia, memory deficits from exposure to anesthetic agents, sleep deprivation induced cognitive impairment, chronic fatigue syndrome, narcolepsy, AIDS-related dementia, epilepsy-related cognitive impairment, Down's syndrome, alcoholism related dementia, drug/substance induced memory impairments, dementia pugilistica (boxer syndrome), or loss of cholinergic synapses. Thus, N-quinuclidinyl-amide I was prepared via an amidation

reaction of 1,2-benzisothiazole-3-carboxylic acid with 3-(R)-aminoquinuclidine dihydrochloride in a 5/1 mixture of THF/DMF using diisopropylethylamine and HATU. $\alpha 7$ NACHR activity of the prepared quinuclidines were determined using rat brain tissue in a competition assay with [3H]-MLA.

AN 2004:287845 CAPLUS

DN 140:321562

TI Preparation of quinuclidinyl indazoles, benzothiazoles and benzoisothiazoles for use in pharmaceutical compositions as nicotinic acetylcholine receptor ligands

IN Tehim, Ashok; Herbert, Brian; Nguyen, Truc Minh; Xie, Wenge; Gauss, Carla Maria

PA Memory Pharmaceuticals Corporation, USA

SO PCT Int. Appl., 147 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004029050	A1	20040408	WO 2003-US29976	20030925
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2499128	AA	20040408	CA 2003-2499128	20030925
	US 2004132790	A1	20040708	US 2003-669645	20030925
	EE 200500011	A	20050615	EE 2005-11	20030925
	EP 1543000	A1	20050622	EP 2003-798723	20030925
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	BR 2003014485	A	20050726	BR 2003-14485	20030925
	JP 2006503851	T2	20060202	JP 2004-540191	20030925
	NO 2005001985	A	20050609	NO 2005-1985	20050422
PRAI	US 2002-413151P	P	20020925		
	US 2003-448469P	P	20030221		
	WO 2003-US29976	W	20030925		

OS MARPAT 140:321562

IT 677306-44-4P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of N-quinuclidinyl indazoles, benzothiazoles and benzoisothiazoles for use in pharmaceutical compns. as nicotinic acetylcholine receptor ligands)

RN 677306-44-4 CAPLUS

CN 5-Benzothiazolecarboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-, monohydrochloride (9CI) (CA INDEX NAME)


Absolute stereochemistry.




● HCl

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 14 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



I



II

AB The present invention is directed to phenylindole based small mol. inhibitors I [L and M independently = H, alkyl, alkoxy, aryl, (un)substituted aryl, hydroxy, halo, amino, alkylamino, etc.; R = H, alkyl, benzyl, 4-fluorobenzyl, and dialkylamino alkyl; R1 and R2 independently = H, (un)substituted-alkyl, -cycloalkyl, -polycyclic aliphatic groups, -Ph, etc.; X = NHCO or CONH] of the IgE response to allergens, which are useful in the treatment of allergy and/or asthma or any diseases where IgE is pathogenic. Methods for preparing intermediates for the synthesis of I are given; thus, e.g., II (Ad = 1-adamantyl) was prepared by condensation of 3-bromophenylhydrazine with 4-nitroacetophenone followed by acid catalyzed cyclocondensation to form 2-(4-nitrophenyl)-6-bromo-1H-indole which underwent substitution from the cyano derivative which was converted in three steps to the 2-(4-nitrophenyl)-1H-indole-6-carboxylic acid, then a sequence of amidation, nitro reduction In assays for determining suppression of IgE

response, I produced 50% inhibition at concentration ranges from 1 pM to 100 μ M. This invention also relates to phenyl-indole mols. that are cellular proliferation inhibitors and thus are useful as anticancer agents. This invention further relates to small mols. which suppress cytokines and leukocytes.

AN 2004:252623 CAPLUS

DN 140:287264

TI Pharmaceutical compositions of phenyl-indole compounds for modulating IgE and inhibiting cellular proliferation

IN Sircar, Jagadish C.; Ramnauth, Jailall; Richards, Mark L.

PA Avanir Pharmaceuticals, USA

SO PCT Int. Appl., 131 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004024896	A2	20040325	WO 2003-US30959	20030912
	WO 2004024896	A3	20040624		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	CA 2498493	AA	20040325	CA 2003-2498493	20030912
	US 2004180946	A1	20040916	US 2003-661139	20030912
	EP 1537079	A2	20050608	EP 2003-795717	20030912
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
	BR 2003014223	A	20050726	BR 2003-14223	20030912
PRAI	US 2002-410777P	P	20020912		
	WO 2003-US30959	W	20030912		

OS MARPAT 140:287264


IT 675822-89-6P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(drug candidate; preparation of phenylindoles as modulators of IgE response and antitumor agents)

RN 675822-89-6 CAPLUS


CN 1H-Indole-5-carboxamide, N-1-azabicyclo[2.2.2]oct-3-yl-2-[4-[(4-methoxybenzoyl)amino]phenyl]- (9CI) (CA INDEX NAME)



L6 ANSWER 15 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



I



II

AB The present invention is directed to phenylindole based small mol. inhibitors I [L and M independently = H, alkyl, alkoxy, aryl, (un)substituted aryl, hydroxy, halo, amino, alkylamino, etc.; R = H, alkyl, benzyl, 4-fluorobenzyl, and dialkylamino alkyl; R1 and R2 independently = H, (un)substituted-alkyl, -cycloalkyl, -polycyclic aliphatic groups, -Ph, etc.; X = NHCO or CONH] of the IgE response to allergens, which are useful in the treatment of allergy and/or asthma or any diseases where IgE is pathogenic. Methods for preparing intermediates for the synthesis of I are given; thus, e.g., II (Ad = 1-adamantyl) was prepared by condensation of 3-bromophenylhydrazine with 4-nitroacetophenone followed by acid catalyzed cyclocondensation to form 2-(4-nitrophenyl)-6-bromo-1H-indole which underwent substitution from the cyano derivative which was converted in three steps to the 2-(4-nitrophenyl)-1H-indole-6-carboxylic acid, then a sequence of amidation, nitro reduction In assays for determining suppression of IgE

response, I produced 50% inhibition at concentration ranges from 1 pM to 100 μ M. This invention also relates to phenyl-indole mols. that are cellular proliferation inhibitors and thus are useful as anticancer agents. This invention further relates to small mols. which suppress cytokines and leukocytes.

AN 2004:252465 CAPLUS

DN 140:287262

TI Pharmaceutical compositions of phenyl-indole compounds for modulating IgE and inhibiting cellular proliferation

IN Sircar, Jagadish C.; Ramnauth, Jailall; Richards, Mark L.

PA Avanir Pharmaceuticals, USA

SO PCT Int. Appl., 112 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

PATENT NO.

KIND

DATE

APPLICATION NO.

DATE

PI WO 2004024655 A2 20040325 WO 2003-US28145 20030909
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
 CO, CR, CU, CZ, CZ, DE, DE, DK, DK, DM, DZ, EC, EE, EE, EG, ES,
 FI, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG,
 KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
 MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
 SK, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
 YU, ZA, ZM, ZW
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG,
 CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC,
 NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
 GW, ML, MR, NE, SN, TD, TG

US 2004180946 A1 20040916 US 2003-661139 20030912
 PRAI US 2002-410777P P 20020912

OS MARPAT 140:287262


IT 675822-89-6P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU
 (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
 (Uses)


(drug candidate; preparation of phenylindoles as modulators of IgE response
 and antitumor agents)

RN 675822-89-6 CAPLUS


CN 1H-Indole-5-carboxamide, N-1-azabicyclo[2.2.2]oct-3-yl-2-[4-[(4-
 methoxybenzoyl)amino]phenyl]- (9CI) (CA INDEX NAME)



L6 ANSWER 16 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 GI



I



II

AB The title compds. [I and II; A, B, C, and D = C, N; W, X, Y and Z = C, N and at least one of W, X, Y, and Z = N; R1-R8 = H, halo, CN, NO₂, etc.; R9 = H, (un)substituted alkyl, aryl, etc.; R10 = H; or NR₉R₁₀ = 5-7 membered ring], useful for inhibiting various enzymes and treating various conditions, were prepared E.g., a multi-step synthesis of 4-amino-3-(benzimidazol-2-yl)-6-(4-methylpiperazinyl)hydroquinolin-2-one, was given. The majority of the exemplary compds. I displayed an IC₅₀ of less than 10 μM with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, Cdk4, MEK1, NEK-2, CHK2, CK1ε, Raf, Fyn, Lck, Rsk2, PAR-1, c-Kit, c-ABL, p60src, FGFR3, FLT-3, PDGFRα, and PDGFRβ. In addition, many of the exemplary compds. exhibited IC₅₀ values in the nM range and show potent activity with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, FGFR3, c-Kit, c-ABL, FLT-3, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, MEK1, CHK2, Fyn, Lck, Rsk2, PAR-1, PDGFRα, and PDGFRβ with IC₅₀ values of less than 1 μM.

AN 2004:182836 CAPLUS

DN 140:235711

TI Preparation of benzimidazole quinolinones for inhibiting a serine/threonine kinase

IN Barsanti, Paul A.; Bussiere, Dirksen; Harrison, Stephen D.; Heise, Carla C.; Jansen, Johanna M.; Jazan, Elisa; Machajewski, Timothy D.; McBride, Christopher; McCrea, William R.; Ng, Simon; Ni, Zhi-Jie; Pecchi, Sabina; Pfister, Keith; Ramurthy, Savithri; Renhowe, Paul A.; Shafer, Cynthia M.; Silver, Joel B.; Wagman, Allan; Weismann, Marion

PA Chiron Corporation, USA

SO PCT Int. Appl., 570 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 7

PATENT NO.

KIND


DATE

APPLICATION NO.

DATE

PI	WO 2004018419	A2	20040304	WO 2003-US25990	20030819
	WO 2004018419	A3	20040603		
	WO 2004018419	B1	20040729		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2496164	AA	20040304	CA 2003-2496164	20030819
	EP 1539754	A2	20050615	EP 2003-781286	20030819
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	BR 2003013743	A	20050705	BR 2003-13743	20030819
	JP 2006503919	T2	20060202	JP 2005-501762	20030819
PRAI	US 2002-405729P	P	20020823		
	US 2002-426107P	P	20021113		
	US 2002-426226P	P	20021113		
	US 2002-426282P	P	20021113		
	US 2002-428210P	P	20021121		
	US 2003-460327P	P	20030403		
	US 2003-460328P	P	20030403		
	US 2003-460493P	P	20030403		
	US 2003-478916P	P	20030616		
	US 2003-484048P	P	20030701		
	WO 2003-US25990	W	20030819		
OS	MARPAT 140:235711				
IT	668429-33-2P				
	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(preparation of benzimidazole quinolinones for inhibiting a serine/threonine kinase)				
RN	668429-33-2 CAPLUS				
CN	1H-Benzimidazole-5-carboxamide, 2-(4-amino-1,2-dihydro-2-oxo-3-quinolinyl)-N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)				


Absolute stereochemistry.




*nevera hel
in applicate*

=> d abs bib fhitstr 1-9

L6 ANSWER 1 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



I



II

AB The title compds. I [A, B, C, and D = C, N; R1-R3 = H, halo, CN, NO2, etc.; R4 = H, alkyl; R5-R8 = H, halo, CN, NO2, etc.; R9 = H, (un)substituted alkyl, aryl, etc.; R10 = H], useful for inhibiting fibroblast growth factor receptor 3 or treating a biol. condition mediated by fibroblast growth factor receptor 3, were prepared E.g., a multi-step synthesis of 4-amino-5-fluoro-3-[6-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-1H-quinolin-2-one (II), starting from 5-chloro-2-nitroaniline and 1-methylpiperazine, was given. The majority of the exemplary compds. I displayed an IC50 of less than 10 μ M with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, Cdk4, MEK1, NEK-2, CHK2, CK1 ϵ , Raf, Fyn, Lck, Rsk2, PAR-1, c-Kit, c-ABL, p60src, FGFR3, FLT-3, PDGFR α , and PDGFR β . In addition, many of the exemplary compds. exhibited IC50 values in the nM range and show potent activity with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, FGFR3, c-Kit, c-ABL, FLT-3, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, MEK1, CHK2, Fyn, Lck, Rsk2, PAR-1, PDGFR α , and PDGFR β with IC50 values of less than 1 μ M. The mentioned above compound II was tested in various tests and showed significant antiproliferative activity. II inhibited FGFR3 receptor phosphorylation and ERK phosphorylation in multiple myeloma cell lines with activating FGFR3 mutations.

AN 2005:1242789 CAPLUS

DN 143:477969

TI Preparation of benzimidazole quinolinones for inhibiting FGFR3 and treating multiple myeloma

IN Cai, Shaopei; Chou, Joyce; Harwood, Eric; Heise, Carla C.; Machajewski,

Timothy D.; Ryckman, David; Shang, Xiao; Wiesmann, Marion; Zhu, Shuguang
 PA Chiron Corporation, USA
 SO U.S. Pat. Appl. Publ., 239 pp., Cont.-in-part of U.S. Ser. No. 644,055.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN. CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2005261307	A1	20051124	US 2004-983174	20041105
	US 2004092535	A1	20040513	US 2003-644055	20030819
	US 2005203101	A1	20050915	US 2004-839793	20040505
PRAI	US 2002-405729P	P	20020823		
	US 2002-426107P	P	20021113		
	US 2002-426226P	P	20021113		
	US 2002-426282P	P	20021113		
	US 2002-428210P	P	20021121		
	US 2003-460327P	P	20030403		
	US 2003-460328P	P	20030403		
	US 2003-460493P	P	20030403		
	US 2003-478916P	P	20030616		
	US 2003-484048P	P	20030701		
	US 2003-644055	A2	20030819		
	US 2003-517915P	P	20031107		
	US 2003-526425P	P	20031202		
	US 2003-526426P	P	20031202		
	US 2004-546017P	P	20040219		

OS MARPAT 143:477969

IT 668429-33-2P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of benzimidazole quinolinones for inhibiting FGFR3 and treating multiple myeloma)


RN 668429-33-2 CAPLUS

CN 1H-Benzimidazole-5-carboxamide, 2-(4-amino-1,2-dihydro-2-oxo-3-quinolinyl)-N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 2 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 GI



AB The title compds. [I; A, B, C, D = C, N; R1 = H, halo, CN, NO₂, etc.; R2, R3 = H, halo, NO₂, CN, etc.; R4 = H, (un)substituted alkyl; R5, R8 = H, (un)substituted alkyl, alkenyl, heterocyclyl; or R5 may be absent if A = N; or R8 may be absent if D = N; R6, R7 = H, halo, NO₂, CN, etc.; R9 = H, (un)substituted alkyl, aryl, etc.; R10 = H; or R9 and R10 join together to form one or more rings, each having 5-7 members], useful for inhibiting checkpoint kinase 1, inducing cell cycle progression, and increasing apoptosis in cells, were prepared E.g., a multi-step synthesis of 4-amino-3-(benzimidazol-2-yl)-6-(4-methylpiperazinyl)hydroquinolin-2-one, was given. The compds. I were tested against various kinases. Two of the prepared compds. I, 4-[(3S)-1-azabicyclo[2.2.2]oct-3-ylamino]-3-(1H-benzimidazol-2-yl)-6-chloroquinolin-2-(1H)-one and 6-chloro-3-[5-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-4-[(piperidin-2-ylmethyl)amino]quinolin-2(1H)-one, were found to be potent inhibitors of CHK1 with IC₅₀ of 0.32 nM and 0.63 nM, resp. The majority of the exemplary compds. I displayed an IC₅₀ of less than 10 μM with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, Cdk4, MEK1, NEK-2, CHK2, CK1ε, Raf, Fyn, Lck, Rsk2, PAR-1, c-Kit, c-ABL, p60src, FGFR3, FLT-3, PDGFRα, and PDGFRβ. In addition, many of the exemplary compds. exhibited IC₅₀ values in the nM range and show potent activity with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, FGFR3, c-Kit, c-ABL, FLT-3, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, MEK1, CHK2, Fyn, Lck, Rsk2, PAR-1, PDGFRα, and PDGFRβ with IC₅₀ values of less than 1 μM. The compds. I may be used to prepare pharmaceutical compns. and may be used in conjunction with DNA damaging agents.

AN 2005:1223876 CAPLUS

DN 143:477966

TI Preparation of benzimidazole quinolinones for inhibiting a checkpoint kinase 1 and their use in combination therapy for cancer

IN Gesner, Thomas G.; Barsanti, Paul A.; Harrison, Stephen D.; Ni, Zhi-Jie; Brammeier, Nathan M.; Zhou, Yasheen; Le, Vincent P.

PA Chiron Corporation, USA

SO U.S. Pat. Appl. Publ., 249 pp., Cont.-in-part of U.S. Ser. No. 644,055. CODEN: USXXCO

DT Patent

LA English

FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2005256157	A1	20051117	US 2005-41191	20050121
	US 2004092535	A1	20040513	US 2003-644055	20030819
	US 2005203101	A1	20050915	US 2004-839793	20040505
PRAI	US 2002-405729P	P	20020823		

US 2002-426107P	P	20021113
US 2002-426226P	P	20021113
US 2002-426282P	P	20021113
US 2002-428210P	P	20021121
US 2003-460327P	P	20030403
US 2003-460328P	P	20030403
US 2003-460493P	P	20030403
US 2003-478916P	P	20030616
US 2003-484048P	P	20030701
US 2003-644055	A2	20030819
US 2004-538984P	P	20040123

OS MARPAT 143:477966

IT 668429-33-2P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of benzimidazole quinolinones for inhibiting a checkpoint kinase 1 and their use in combination therapy for cancer)


RN 668429-33-2 CAPLUS

CN 1H-Benzimidazole-5-carboxamide, 2-(4-amino-1,2-dihydro-2-oxo-3-quinolinyl)-N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 3 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI




AB The invention provides methods for using of using 4-Amino-3-(benzimidazol-2-yl)quinolin-2-one derivs. (Markush included), or a salt or tautomer thereof, in the treatment of disorders relating to cell adhesion and metastatic processes. Preparation of I is included.

AN 2005:976928 CAPLUS
 DN 143:279443
 TI 4-Amino-3-(benzimidazol-2-yl)quinolin-2-one derivatives for the modulation of inflammatory and metastatic processes
 IN Lee, Sang H.; Heise, Carla C.
 PA Chiron Corporation, USA
 SO PCT Int. Appl., 145 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005082340	A2	20050909	WO 2005-US5316	20050218
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	US 2005239825	A1	20051027	US 2005-61386	20050218
PRAI	US 2004-546395P	P	20040220		
	US 2004-547103P	P	20040223		
	US 2004-554771P	P	20040319		
OS	MARPAT 143:279443				
IT	668429-33-2				
	RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (benzimidazolyl aminoquinolinone derivs. for modulation of inflammatory and metastatic processes)				
RN	668429-33-2 CAPLUS				
CN	1H-Benzimidazole-5-carboxamide, 2-(4-amino-1,2-dihydro-2-oxo-3-quinolinyl)-N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)				

Absolute stereochemistry.



L6 ANSWER 4 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 AB This invention relates to combinations of an atypical antipsychotic, and a nicotinic receptor agonist or antagonist, kits containing such combinations,


pharmaceutical compns. comprising such combinations, and methods of using such combinations to treat patients suffering from cognitive impairment disorders or psychotic disorders or conditions. A composition was prepared by combining ziprasidone with the nicotinic agonist varenicline tartrate.

AN 2005:612120 CAPLUS
 DN 143:139163
 TI Combination of an atypical antipsychotic and a nicotinic receptor agonist or antagonist for cognition enhancement and psychotic disorders
 IN Romano, Steven Joseph
 PA Pfizer Products Inc., USA
 SO PCT Int. Appl., 62 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1



	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005063296	A2	20050714	WO 2004-IB4174	20041215
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	US 2005215571	A1	20050929	US 2004-18100	20041220
PRAI	US 2003-532082P	P	20031223		
OS	MARPAT 143:139163				
IT	478148-80-0				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (combination of an atypical antipsychotic and a nicotinic receptor agonist or antagonist for cognition enhancement and psychotic disorders)				
RN	478148-80-0 CAPLUS				
CN	Furo[2,3-c]pyridine-5-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-3-methyl- (9CI) (CA INDEX NAME)				

not good

Absolute stereochemistry.



L6 ANSWER 5 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 GI



AB The title compds. I [A, B, C, and D = C, N; R1-R3 = H, halo, CN, NO2, etc.; R4 = H, alkyl; R5-R8 = H, halo, CN, NO2, etc.; R9 = H, (un)substituted alkyl, aryl, etc.; R10 = H], useful for inhibiting fibroblast growth factor receptor 3 or treating a biol. condition mediated by fibroblast growth factor receptor 3, were prepared E.g., a multi-step synthesis of 4-amino-5-fluoro-3-[6-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-1H-quinolin-2-one (II), starting from 5-chloro-2-nitroaniline and 1-methylpiperazine, was given. The majority of the exemplary compds. I displayed an IC50 of less than 10 μ M with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, Cdk4, MEK1, NEK-2, CHK2, CK1 ϵ , Raf, Fyn, Lck, Rsk2, PAR-1, c-Kit, c-ABL, p60src, FGFR3, FLT-3, PDGFR α , and PDGFR β . In addition, many of the exemplary compds. exhibited IC50 values in the nM range and show potent activity with respect to VEGFR1, VEGFR2, VEGFR3, FGFR1, FGFR3, c-Kit, c-ABL, FLT-3, CHK1, Cdc2, GSK-3, NEK-2, Cdk2, MEK1, CHK2, Fyn, Lck, Rsk2, PAR-1, PDGFR α , and PDGFR β with IC50 values of less than 1 μ M. The mentioned above compound II was tested in various tests and showed significant antiproliferative activity. II inhibits FGFR3 receptor phosphorylation and ERK phosphorylation in multiple myeloma cell lines with activating FGFR3 mutations.

AN 2005:451351 CAPLUS

DN 143:7710

TI Preparation of benzimidazole quinolinones for inhibiting FGFR3 and treating multiple myeloma

IN Cai, Shaopei; Chou, Joyce; Harwood, Eric; Heise, Carla C.; Machajewski, Timothy D.; Ryckman, David; Shang, Xiao; Wiesmann, Marion; Zhu, Shuguang

PA Chiron Corporation, USA

SO PCT Int. Appl., 567 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 7

PATENT NO.

KIND

DATE

APPLICATION NO.

DATE

PI WO 2005047244 A2 20050526 WO 2004-US36956 20041105

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2005137399 A1 20050623 US 2004-982757 20041105

US 2005209247 A1 20050922 US 2004-982543 20041105

PRAI US 2003-517915P P 20031107

US 2003-526425P P 20031202

US 2003-526426P P 20031202

US 2004-546017P P 20040219

OS MARPAT 143:7710

IT 668429-33-2P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(preparation of benzimidazole quinolinones for inhibiting FGFR3 and treating multiple myeloma)

RN 668429-33-2 CAPLUS

CN 1H-Benzimidazole-5-carboxamide, 2-(4-amino-1,2-dihydro-2-oxo-3-quinolinyl)-N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)


date is good

Absolute stereochemistry.




L6 ANSWER 6 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

GI



I



II

AB The title compds. I [R1-R4 = H, halo, CN, NO2, etc.; R5-R8 = H, halo, NO2, etc.; R9 = H; R12 = H, alkyl, aryl, heterocyclyl; R13 = H, alkyl, aryl, heterocyclyl, etc.; R14 = H] and their pharmaceutically acceptable lactate salts, useful for inhibiting vascular endothelial growth factor receptor tyrosine kinase, were prepared E.g., a multi-step synthesis of 4-amino-5-fluoro-3-[6-(4-methylpiperazin-1-yl)-1H-benzimidazol-2-yl]-1H-quinolin-2-one (II) and its lactate salt, starting from 5-chloro-2-nitroaniline and 1-methylpiperazine, was given. The pharmaceutically acceptable salts of I have improved aqueous solubility and desirable drug substance properties. Many of the exemplary compds. I displayed an IC50 of less than 10 μ M with respect to Flt-1, KDR, PDGF, c-KIT, FLT-3, VEGFR1, VEGFR2, c-Met, CSF-1, FGFR3 and/or bFGFR. In addition, many of the exemplary compds. exhibited IC50 value of less than 10 μ M with respect to PDGFR. The 4-amino substituted compds. I such as II were found to be potent inhibitors of various kinases such as VEGFR2 (KDR, Flk-1), FGFR1 and PDGFR β with IC50's ranging from 10-27 nM. II inhibits FGFR3 receptor phosphorylation and ERK phosphorylation in multiple myeloma cell lines with activating FGFR3 mutations.

AN 2005:451118 CAPLUS

DN 143:7709

TI Preparation of benzimidazole quinolinones and lactate salts thereof for inhibiting vascular endothelial growth factor receptor tyrosine kinase

IN Cai, Shaopei; Chou, Joyce; Harwood, Eric; Machajewski, Timothy D.; Ryckman, David; Shang, Xiao; Zhu, Shuguang

PA Chiron Corporation, USA

SO PCT Int. Appl., 215 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005046589	A2	20050526	WO 2004-US36941	20041105
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,				

CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

US 2005137399 A1 20050623 US 2004-982757 20041105
 US 2005209247 A1 20050922 US 2004-982543 20041105

PRAI US 2003-517915P P 20031107
 US 2003-526425P P 20031202
 US 2003-526426P P 20031202
 US 2004-546017P P 20040219

OS MARPAT 143:7709

IT 668429-33-2P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of benzimidazole quinolinones and lactate salts thereof for inhibiting vascular endothelial growth factor receptor tyrosine kinase)

RN 668429-33-2 CAPLUS

CN 1H-Benzimidazole-5-carboxamide, 2-(4-amino-1,2-dihydro-2-oxo-3-quinolinyl)-N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L6 ANSWER 7 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AB The present invention relates to compositions and methods to treat diseases or conditions with alpha-7 nicotinic acetylcholine receptor (AChR) full agonists by decreasing levels of tumor necrosis factor-alpha and/or by stimulating vascular angiogenesis.

AN 2004:633526 CAPLUS

DN 141:167817

TI Treatment of diseases with alpha-7 NACH receptor full agonists

IN Groppi, Vincent Edward, Jr.; Rogers, Bruce Nelsen; Rudmann, Daniel Gregory

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 142 pp.

CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004064836	A2	20040805	WO 2004-IB115	20040112
	WO 2004064836	A3	20041223		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ				
	CA 2513433	AA	20040805	CA 2004-2513433	20040112
	EP 1587511	A2	20051026	EP 2004-701414	20040112
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	US 2006019984	A1	20060126	US 2004-761914	20040121
PRAI	US 2003-441801P	P	20030122		
	WO 2004-IB115	W	20040112		
OS	MARPAT 141:167817				
IT	478148-80-0P, N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-3-methylfuro[2,3-c]pyridine-5-carboxamide				
	RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(nAChR agonist; preparation of N-(quinuclidinyl)heteroarylamides as nAChR agonists for use in combination therapy for treatment of ADHD)				
RN	478148-80-0 CAPLUS				
CN	Furo[2,3-c]pyridine-5-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-3-methyl- (9CI) (CA INDEX NAME)				

done not good

Absolute stereochemistry.



L6 ANSWER 8 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



I

AB The invention provides a preparation of mono and hemi fumarate salts of furo[2,3-c]pyridine derivative I, useful as $\alpha 7$ nAChR agonist. The obtained fumarate salts are useful to treat diseases such as Alzheimer's disease, schizophrenia, and arteriosclerosis, etc. (no biol. data). The

title compound I•HO₂CC:CCO₂H was prepared via amidation of furo[2,3-c]pyridine-5-carboxylic acid with (R)-3-aminoquinuclidine and subsequent fumarate salt formation.

AN 2004:515515 CAPLUS

DN 141:54316

TI A preparation of crystalline fumarate salts of furo[2,3-c]pyridine derivative, useful as α 7 nAChR agonists

IN Selbo, Jon Gordon; Hewitt, Bradley Dee; Rappath, David Warner; Wishka, Donn Gregory; Sheikh, Ahmad Yahya

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 46 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1


	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004052894	A1	20040624	WO 2003-IB5607	20031201
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2506529	AA	20040624	CA 2003-2506529	20031201
	EP 1572700	A1	20050914	EP 2003-812627	20031201
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	BR 2003017019	A	20051025	BR 2003-17019	20031201
	US 2005165047	A1	20050728	US 2004-962071	20041008
	NO 2005002560	A	20050817	NO 2005-2560	20050526
PRAI	US 2002-431619P	P	20021206		
	WO 2003-IB5607	W	20031201		
	US 2003-729286	B1	20031205		
IT	708261-37-4P				
	RL: PAC (Pharmacological activity); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(crystal structure; preparation of crystalline fumarate salts of (azabicyclooctyl)furopyridine derivative, useful as α 7 nAChR agonists)				
RN	708261-37-4 CAPLUS				
CN	Furo[2,3-c]pyridine-5-carboxamide, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl-, (2E)-2-butenedioate (1:1) (9CI) (CA INDEX NAME)				

CM 1

CRN 478149-53-0

CMF C15 H17 N3 O2

Absolute stereochemistry.




CM 2


CRN 110-17-8

CMF C4 H4 O4

Double bond geometry as shown.



L6 ANSWER 9 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
GI



AB The compds. of formula: azabicyclo-NH-CO-W [azabicyclo = (substituted) [2.2.2], [2.2.1] or [3.2.1] azabicyclo ring; W = (substituted) pyrazolyl, benzofuranyl, furopyridyl, thiophenyl, furyl, Ph, etc.] as prepared, and are labeled with a radioactive isotopic moiety such as ^{11}C , ^{18}F , ^{76}Br , ^{123}I or ^{125}I . Radiolabeled ligands useful as probes for determining the relative abundance, receptor occupancy, and/or function of nicotinic acetylcholine receptors. Disorders are diagnosed by administering to a mammal a detectably labeled compound and detecting the binding of that compound to the nAChR. The compds. that have been administered are detected using methods including, but not limited to, position emission tomog. and single-photon to emission computed tomog. The present invention is useful in diagnosing a wide variety of diseases and disorders as discussed herein. Thus, I was prepared from Ph chloroformate, 4-iodopyrazole and (R)-(+)-3-aminoquinuclidine dihydrochloride.

AN 2004:515511 CAPLUS

DN 141:54518

TI Preparation of N-azabicyclo carboxamides as radioligands for the diagnosis of disease

IN Rogers, Bruce Nelsen; Piotrowski, David Walter; Groppi, Vincent Edward, Jr.; Jacobsen, Eric Jon; Myers, Jason Kenneth; Walker, Daniel Patrick; Wishka, Donn Gregory; Skaddan, Marc Bradley

PA Pharmacia & Upjohn Company, USA

SO PCT Int. Appl., 80 pp.

CODEN: PIXXD2

DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004052889	A1	20040624	WO 2003-IB5521	20031124
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE,				
	GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,				
	LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ,				
	OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,				
	TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,				
	BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE,				
	ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,				
	TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	US 2004157878	A1	20040812	US 2003-729529	20031205
PRAI	US 2002-431473P	P	20021206		
OS	MARPAT 141:54518				
IT	706782-57-2P				
	RL: DGN (Diagnostic use); SPM (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)				
	(preparation of N-azabicyclo carboxamides as radioligands for diagnosis of disease)				
RN	706782-57-2 CAPLUS				
CN	5-Benzofurancarboxamide-11C, N-(3R)-1-azabicyclo[2.2.2]oct-3-yl- (9CI)				
	(CA INDEX NAME)				

Absolute stereochemistry.

